

INITIAL SOLAR IRRADIANCE MEASUREMENTS FROM THE NOAA-14 SBUV/2 INSTRUMENT

E. Hilsenrath

NASA Goddard Space Flight Center

M. T. DeLand¹ and R. P. Cebula¹

Hughes STX Corporation

¹Supported by NASA Grant NASW-4864 and NASA Contract NAS5-31755

EOS Trans. Amer. Geophys. Union, 76(17), Spring Meeting Suppl. S239, 1995


(NASA-CR-200304) INITIAL SOLAR
IRRADIANCE MEASUREMENTS FROM THE
NOAA-14 SBUV/2 INSTRUMENT (NASA.
Goddard Space Flight Center) 10 p

N96-18510

Unclas

NASA
IN-46
8171
P-10

SPACE BORNE SOLAR UV IRRADIANCE OBSERVATION



SBUV/2 ON NOAA POLAR ORBITING SATELLITES FOR OZONE AND SOLAR IRRADIANCE MEASUREMENTS

1/4 meter double-beam spectrometer

1.1 nm resolution

160 - 405 nm

Solar Views

Sweep mode (full spectrum includes Mg II at 0.15 nm) - once per day

Discrete mode (12 steps around Mg II line) - once per day


Discrete mode (ozone channels) - once per day

Absolute calibration is not internally tracked




Wavelength calibration is tracked to <0.02 nm

Mg index does not require absolute radiometric calibration


NOAA-14 Absolute Irradiance for 1995/ 41




NOAA-14 SBUV/2 Irradiance Comparisons




SBUV-series Mg II INDEX DATA: 27 day RUNNING average



COMPOSITE Mg II SOLAR UV ACTIVITY INDEX






L

L

Solar Activity Measurements



CONCLUSIONS

NOAA-14, beginning in February 1995, continues series of SBUV/2 solar irradiance measurements which began in 1984.

NOAA-11 terminated solar observations in October 1994.

NOAA-9 continues to operate and can provide overlap with considerable work.

Initial results from NOAA-14

Calibration offset between SUSIM and SSBUV

Grating drive unlock problems

Show Mg II variations are detectable during solar minimum with 13 day variations consisted with 10.7 cm solar flux